

# Alcohol Use Disorder in Youth: Information for Physicians



Image credit: Adobe Stock

**Summary:** Many youth will experiment with alcohol and substances. Some will continue to have problems with alcohol and drug abuse. Primary care physicians can help identify and screen youth for problems such as by using the CRAAFT. Using strategies such as motivational interviewing, primary care physicians can also directly help youth to stop their alcohol use, as well as help direct them to specialized addictions treatment.

## Epidemiology

- 10% of the population abuses drugs/alcohol, and 20% of patients seen by family physicians have substance-abuse problems (not including tobacco) (Mersy, 2003)
- Prevalence of DSM-IV alcohol abuse and alcohol dependence in the US between 2001 and 2002 was estimated to be
  - Alcohol abuse — 17.8 percent lifetime, 4.7 percent past 12 months
  - Alcohol dependence — 12.5 percent lifetime, 3.8 percent past 12 months (Hassin, 2011)

## Signs/Symptoms

### Medical problems

- Frequent trauma or accidents
- GI problems such as epigastric distress, diarrhea, weight changes
- Sexual problems

### Psychiatric

- Mood or anxiety problems

### Impaired function

- Decreased social function
- Changes in friends, with decreased involvement with old friends and more involvement with new friends who use drugs, along with reluctance to introduce parents to new friends
- Decreased function at home
- Decreased function at school
- Skipping classes and missing days

- Worsening grades
- Decreased function at work
- Frequent absences

## Screening

---

CRAFFT is a mnemonic for screening addictions for children/youth up to age 21, recommended by the American Academy of Paediatrics.

- C)ar: "Have you ever ridden in a CAR driven by someone (including yourself) who was "high" or had been using alcohol or drugs?"
- R)elax: "Do you ever use alcohol or drugs to RELAX, feel better about yourself, or fit in?"
- A)lone: "Do you ever use alcohol/drugs while you are by yourself, ALONE?"
- F)orget: "Do you ever FORGET things you did while using alcohol or drugs?"
- F)riends: "Do your family or FRIENDS ever tell you that you should cut down on your drinking or drug use?"
- T)rouble: "Have you gotten into TROUBLE while you were using alcohol or drugs?"

## DSM-5

---

### Criteria for Alcohol Abuse

1. Alcohol is often taken in larger amounts or over a longer period than was intended.
2. There is a persistent desire or unsuccessful efforts to cut down or control alcohol use.
3. A great deal of time is spent in activities necessary to obtain alcohol, use alcohol or recover from its effects.
4. Craving or a strong desire or urge to use alcohol.
5. Recurrent alcohol use resulting in a failure to fulfill major role obligations at work, school or home.
6. Continued alcohol use despite having recurrent social or interpersonal problems.
7. Social, occupational or recreational activities are given up or reduced because of alcohol use.
8. Recurrent alcohol use in situations in which it is physically hazardous.
9. Alcohol use is continued despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by alcohol.
10. Tolerance, as defined by either of the following:
  - a) Increased amounts of alcohol are needed to achieve the same effect
  - b) Marked diminished effect with use of the same amount of alcohol
11. Withdrawal, as manifested by either of the following:
  - a) The characteristic withdrawal syndrome for alcohol
  - b) Alcohol (or a closely related substance such as a benzodiazepine) is taken to relieve or avoid withdrawal symptoms

### DSM 5 criteria for Alcohol Intoxication

- Recent use of substance
- Clinically significant maladaptive behavioural or psychological changes that developed during, or shortly after use (inappropriate sexual or aggressive behaviour, mood lability, impaired judgement)
- One (or more) of the following signs or symptoms developing during, or shortly after use
  - Slurred speech
  - Incoordination
  - Unsteady gait
  - Nystagmus
  - Impairment in cognition (e.g. attention, memory)
  - Stupor or coma
- Signs or symptoms not attributable to another medical condition, mental disorder, including intoxication with another substance.

[More...](#)

## DDx

---

- Mood and anxiety disorders can both be a result of drug use, but also, drug use can be an attempt to cope with or self-medicate a mood and anxiety disorders
- Attention deficit hyperactivity disorder (ADHD)

## Physical Exam (Px)

---

- Signs of Alcohol Abuse
  - Mild tremor
  - Odor of alcohol on breath
  - Enlarged, tender liver
  - “Aftershave/mouthwash” syndrome (to mask the odor of alcohol)
  - Labile blood pressure, tachycardia (suggestive of alcohol withdrawal)
  - Signs of Liver failure
    - Jaundice
    - Scleral icterus
    - Palmar erythema
    - Spider angioma
    - Caput medusa
    - Ascites
    - Easy bruising
    - Gynecomastia
    - Asterixis
    - Etc.
- Signs of concurrent substance abuse
  - Nasal irritation (suggestive of cocaine insufflation)
  - Conjunctival irritation (suggestive of exposure to marijuana smoke)
  - Odor of marijuana on clothing

## Investigations

---

- Routine drug screening in the primary care office is not recommended, unless it is part of a comprehensive plan, generally working in conjunction with other substance use or mental health professionals
- Common investigations include:
  - Urine drug screening for marijuana use
  - CBC (anemia, thrombocytopenia)
  - GGT, MCV, carbohydrate-deficient transferrin for alcohol use
  - Liver indices for liver impairment
    - AST:ALT approaches two with alcoholism
    - INR (decreased clotting factor production by liver)
  - BUN/CR for renal impairment

## Management

---

- Work with Families and Caregivers

Validate whatever support the family/caregiver has provided, e.g. Clinician: "I am so grateful that you have been a support for your (loved one). If not for your support, I suspect things would be even worse."

- Encourage families and carers to be involved; the individual with alcohol use needs the support of other family members and carers, e.g. Clinician: "Your ongoing support will make a big difference for your family member."
  - Ask about family's need for support, e.g. "How can I support you, so that you can support your loved one?"
  - Provide written and verbal information on alcohol misuse and its management, including how families and carers can support the service user.
- Psychotherapy
    - Approaches depending on the clinical situation may include motivational enhancement therapy, CBT, marital and family therapy
  - Office-based counseling using motivational interviewing techniques
    - Relate patient's presenting problem to the alcohol/substance use
      - E.g. Physician: "You have mentioned that you are having problems with low energy, and troubles at school and work. I wonder if this might have something to do with your alcohol use. What do you think?"

[More...](#)

## Medication Management of Alcohol Abuse

- Medications for withdrawal include
  - Diazepam (Valium) for alcohol withdrawal
    - Start at 5-20 mg po q 1-4 hr up to three times daily
    - If patient is delirious, then start 60 mg immediately
  - Chlordiazepoxide (Librium) for alcohol withdrawal
    - Start at 15-100 mg po daily, max 300 mg daily
- Medications for recovery include
  - Disulfiram (Antabuse), supervised oral disulfiram may be used to prevent relapse but patients must be informed that this is a treatment requiring complete abstinence and be clear about the dangers of taking alcohol with it.
    - After 72-hrs of being alcohol free, start at 125 mg po q AM, then increase to 250 mg daily, max 500 mg daily
    - Labs: Order hepatic profile at baseline, then q monthly x 2-months, then q 3-6 months
  - Naltrexone (ReVia)
    - Start at 25 mg po daily x 2-days, then if no withdrawal symptoms, increase to 50 mg daily
    - Labs: Urine opioid screen, CBC, hepatic profile (AST, ALT, GGT, bilirubin), periodic hepatitis screen
  - Acamprosate (Campral), recommended in newly detoxified dependent patients as an adjunct to psychosocial interventions
    - Target dosage 666 mg daily three times a day
  - Topiramate (Topamax)
- If Wernicke-Korsakoff syndrome
  - Thiamine 100 mg po (or IM) x 3-days (to be given before the patient resumes eating well)
- If depressive symptoms persist for more than two weeks following treatment for alcohol dependence, consider
  - Selective serotonin reuptake inhibitor (SSRI) for mood/anxiety problems
  -

- Bupropion SR (Zyban / Wellbutrin) for smoking cessation
  - Start at 150 mg daily x 2-4 days, then increase to 150 mg po bid, max 300-450 mg divided bid
  - Monitor after a few weeks

## Reporting Obligations

---

For driving: In most jurisdictions, the law requires that physicians report patients who, in the opinion of the physician, may be unfit to drive for medical reasons.

View CMA website for [links to provincial and territorial forms..](#)

[More...](#)

For parenting: If a physician is concerned that alcohol/drug use may be affecting their ability to parent their child(ren), then the law requires that physicians report patients who, in the opinion of the physician, may be neglectful (or abusive) of their children.

## When to Refer

---

- For moderate to severe problems, or problems with alcohol/substance dependence, refer to addictions services
  - Involve the patient in the process
  - Discuss various options
  - Make the appointment for the addiction service with the patient, during the office visit
- If there are alcohol use issues plus mental health symptoms such as depression that persist for more than two weeks following treatment for alcohol dependence, consider referring to addictions / mental health services

## Who to Refer to

---

- Self-help groups
  - Alcoholics Anonymous: In general, groups are open to all ages, and depending on the location, there may also be specific youth groups available as well.
- Alcohol and substance abuse treatment centre

### 1. The following is a good screening measure for alcohol use disorder in youth:

- CRAAFT
- CAGE
- PHQ-9
- AUDIT-10

### 2. Alcohol abuse can be managed in the following ways EXCEPT:

- Psychotherapy
- Working with families and caregivers
- Using disulfiram (Antabuse) in a patient consuming moderate quantities of alcohol.
- Using diazepam (Valium) for alcohol withdrawal.

## References

---

- Mersy D: Recognition of Alcohol and Substance Abuse, *Am Fam Physician*, 2003 Apr 1;67(7):1529-32. Retrieved Sep 9, 2012 from <http://www.aafp.org/afp/2003/0401/p1529.html>
- Scottish Intercollegiate Guidelines Network (SIGN). The management of harmful drinking and alcohol dependence in primary care. A national clinical guideline. Edinburgh (Scotland): Scottish Intercollegiate Guidelines Network (SIGN); 2003 Sep. 39 p. (SIGN publication; no. 74). [158 references]
- National Institute for Health and Clinical Excellence (NICE). Alcohol-use disorders. Diagnosis, assessment and management of harmful drinking and alcohol dependence. London (UK): National Institute for Health and Clinical Excellence (NICE); 2011 Feb. 54 p. (Clinical guideline; no. 115).
- Work Group on Substance Use Disorders, Kleber et al.: Treatment of patients with substance use disorders, second edition. American Psychiatric Association. *Am J Psychiatry* 2006 Aug;163(8 Suppl):5-82. <http://guideline.gov/content.aspx?id=9316&search=alcohol+use>
- Lussier MT, Richard C. The Motivational Interview. *Can Fam Physician*. 2007 December; 53(12);2117-2118. Retrieved Sep 7, 2012 from <http://www.cfp.ca/content/53/12/2117.full>
- Miller RW, Rollnick S. Motivational interviewing. New York, NY: The Guilford Press; 1991.
- Physician's Guide to Helping Patients with Alcohol Problems, NIMH
- Willenbring M, Massey S, Gardner M: *Am Fam Physician*. 2009 Jul 1;80(1):44-50. Retrieved Sep 7, 2012 from <http://www.aafp.org/afp/2009/0701/p44.html>
- Zimmerman G et al.: A 'Stages of Change' Approach to Helping Patients Change Behaviour, *American Family Physician*. 2000 Mar 1;61(5):1409-1416, Retrieved Sep 9, 2012 from <http://www.aafp.org/afp/2000/0301/p1409.html>

---

## About this Document

Written by Dr's Michael Cheng, FRCP(C), Psychiatrist, University of Ottawa, and Dr. M. St-Jean, MD, CCFP, University of Ottawa. Special thanks to Alexandra Birk-Urovitz (uOttawa Class of Medicine) and Elizabeth Birk-Urovitz (family physician) for help with multiple choice questions.

---

## Conflicts of Interest

eMentalHealth.ca/PrimaryCare has received an unrestricted educational grant for eMentalHealth.ca/PrimaryCare from Lundbeck/Otsuka, which markets Citalopram (Celexa). In order to mitigate the potential for bias, Lundbeck/Otsuka are not involved in content development in any way.

---

## Disclaimer

This information is offered 'as is' and is meant only to provide general information that supplements, but does not replace the information from a qualified expert or health professional. Always contact a qualified expert or health professional for further information in your specific situation or circumstance.

---

## Creative Commons License

You are free to copy and distribute this material in its entirety as long as 1) this material is not used in any way that suggests we endorse you or your use of the material, 2) this material is not used for commercial purposes (non-commercial), 3) this material is not altered in any way (no derivative works). View full license at <http://creativecommons.org/licenses/by-nc-nd/2.5/ca/>

---

## Authors

Written by the eMentalHealth Team and Partners. Information partners include members of the Division of Child Psychiatry as well as members of the Department of Family Medicine at the University of Ottawa.